

---

**Articular cartilage regeneration by activated skeletal stem cells.**

**Journal:** Nat Med

**Publication Year:** 2020

**Authors:** Matthew P Murphy, Lauren S Koepke, Michael T Lopez, Xinming Tong, Thomas H Ambrosi, Gunsagar S Gulati, Owen Marecic, Yuting Wang, Ryan C Ransom, Malachia Y Hoover, Holly Steininger, Liming Zhao, Marcin P Walkiewicz, Natalina Quarto, Benjamin Levi, Derrick C Wan, Irving L Weissman, Stuart B Goodman, Fan Yang, Michael T Longaker, Charles K F Chan

**PubMed link:** 32807933

**Funding Grants:** Enhancing healing via Wnt-protein mediated activation of endogenous stem cells

**Public Summary:**

**Scientific Abstract:**

Osteoarthritis (OA) is a degenerative disease resulting in irreversible, progressive destruction of articular cartilage(1). The etiology of OA is complex and involves a variety of factors, including genetic predisposition, acute injury and chronic inflammation(2-4). Here we investigate the ability of resident skeletal stem-cell (SSC) populations to regenerate cartilage in relation to age, a possible contributor to the development of osteoarthritis(5-7). We demonstrate that aging is associated with progressive loss of SSCs and diminished chondrogenesis in the joints of both mice and humans. However, a local expansion of SSCs could still be triggered in the chondral surface of adult limb joints in mice by stimulating a regenerative response using microfracture (MF) surgery. Although MF-activated SSCs tended to form fibrous tissues, localized co-delivery of BMP2 and soluble VEGFR1 (sVEGFR1), a VEGF receptor antagonist, in a hydrogel skewed differentiation of MF-activated SSCs toward articular cartilage. These data indicate that following MF, a resident stem-cell population can be induced to generate cartilage for treatment of localized chondral disease in OA.

---

**Source URL:** <https://www.cirm.ca.gov/about-cirm/publications/articular-cartilage-regeneration-activated-skeletal-stem-cells>